

14% BROWN BAG SPECIAL 3/16"

341

ALL PURPOSE PELLET FOR ALL CLASSES OF CATTLE
ON PASTURE, AND MATURE HORSES

GUARANTEED ANALYSIS

Crude Protein – (min)	14.0%
Crude Fat – (min)	6.0%
Crude Fiber – (max)	20.0%
ADF – (max)	23.0%
Calcium – (min)	2.0%
Calcium – (min)	3.0%
Phosphorus – (min)	0.7%
Potassium – (min)	0.7%
Salt – (min)	0.7%
Salt – (max)	1.7%
Copper – (min)	20 ppm
Selenium – (min)	0.4 ppm
Zinc – (min)	110 ppm
Vitamin A – (min)	10,000 IU/LB
Vitamin D – (min)	1000 IU/LB
Vitamin E – (min)	12 IU/LB

INGREDIENT STATEMENT

Processed Grain By-Products, Roughage Products 25%, Plant Protein Product, Calcium Carbonate, Sodium Bentonite, Salt, Magnesium Oxide, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Vegetable Oil, Sodium Selenite, Cobalt Sulfate, Ethylenediamine Dihydriodide, Vitamin A Supplement, Vitamin D₃ Supplement, Vitamin E Supplement, Propionic Acid (as a preservative).

FEEDING DIRECTIONS

GENERAL RECOMMENDATIONS FOR ALL ANIMALS INCLUDE THESE GUIDELINES. Do not feed more than 1% of body weight and do not to exceed 8 lb in one feeding (the exception would be creep feeding as provided below). Provide fresh water, minerals with salt and good quality hay or forage at all times.

FOR CATTLE UP TO 400 POUNDS – Feed as a creep feed free choice. **THE TWO MOST IMPORTANT THINGS ABOUT CREEP FEEDING ARE TO KEEP THE FEED FRESH AND ALWAYS HAVE FEED AVAILABLE.** Start feeding creep feed at 60-90 days of age. Place the creep feeder in social centers such as watering areas, feeding areas, where the mineral feeder is located, etc. Set the door on the feeder to where there is some feed available, but not an excess. When starting a group of calves on creep feed, start out with a small amount of feed first until they are used to eating the feed and coming to the feeder regularly. Small calves will only eat a small amount per day at first; generally 0.5-1.0 pound per day to start is all they will eat. **ALTERNATIVELY** this feed may be fed daily up

to 1% of body weight per feeding and up to 3 equal feedings per day separated by a minimum of 6 hours.

FOR CATTLE FROM 400 TO 800 POUNDS – This feed may be fed as a creep feed **IF** the cattle are already eating it as a creep feed as described above. **IF NOT – THEY MAY BE FED THIS FEED AS A CREEP FEED AFTER AN ADJUSTMENT PERIOD OF INCREASING FEED INTAKE.** The procedure normally recommended for acclimatization of calves to feed is to feed the calves twice a day (as a minimum) starting at about ½ pound per 100 pounds of body weight. Then up the amount of feed fed by ½ a pound of feed per head per day until feed is being left from one feeding to the other. This should be at about 2-2 1/2 % of body weight. At this time feed may be placed in a feeder and be given free choice to the calves. **ALTERNATIVELY** this feed may be fed daily up to 1% of body weight per feeding and up to 3 equal feedings per day separated by a minimum of 6 hours. Do not feed more than 1% of body weight and do not to exceed 8 lb in one feeding.

FOR BREEDING AGE CATTLE AND COWS (NURSING OR DRY) – Feed from 0.5% up to 2.0% of body weight and do not to exceed 8 lb in one feeding. In times of high stress and poor forage quality or availability, feed more than one time per day. Separate feedings by a minimum of 6 hours for best results.

FOR BREEDING BULLS – Feed at the rate of 0.5 to 1.5% of body weight per head per day depending on forage availability and quality. Do not to exceed 8 lb in one feeding. In times of high work load and poor forage quality or availability, feed more than one time per day. Separate feedings by a minimum of 6 hours for best results.

FOR MATURE HORSES - Do not feed or water extremely hot, tired animals. Feed at the rate of 0.5-1.5 lb. of feed per 100 lbs. of body weight for mature horses. Split feedings into at least two feedings per day if over 5 pounds of feed is fed per day. Feeding levels over 10 pounds per head per day should be done in three split feedings. **DO NOT FEED OVER 5 POUNDS PER 1000 POUNDS OF BODY WEIGHT IN ANY ONE FEEDING.**

Warning: Contains supplemental Copper. Do not feed to Sheep.

Manufactured by
SABINE VALLEY FEEDS
3001 Bois d'Arc
Greenville, Texas 75401

Net Wt. 50 lbs. (22.6 Kg)