

SABINE VALLEY FORTIFIED RANGE MEAL with CTC

#414

For Beef Cattle on Pasture

MEDICATED FEED

For control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella spp.* susceptible to Chlortetracycline.

ACTIVE DRUG INGREDIENT

Chlortetracycline..... 270 gm/ton

- Consumption of approximately 2.6 pounds per 1000 pound animal daily helps prevent some forms of shipping fever.
- Supplies the vitamins and minerals normally consumed with a high quality breeder mineral
- Consumption of approximately 3 pounds supplies the energy equivalent of 2 pounds of corn

GUARANTEED ANALYSIS

Crude Protein – (min)	14.0%
Crude Fat – (min)	8.5%
Crude Fiber – (max).....	8.0%
Calcium – (min)	2.6%
Calcium – (max).....	3.6%
Phosphorus – (min).....	1.0%
Salt – (min)	24.0%
Salt – (max)	29.0%
Potassium - (min)	0.60%
Manganese – (min)	375 PPM
Copper – (min)	80 PPM
Zinc – (min)	375 PPM
Selenium – (min)	1.5 PPM
Cobalt – (min)	4.0 PPM
Iodine – (min)	5.0 PPM
Vitamin A – (min)	20,000 IU/LB
Vitamin D – (min)	5,000 IU/LB
Vitamin E – (min)	20 IU/LB

INGREDIENT STATEMENT

Processed Grain By-Products, Salt, Plant Protein Products, Molasses Products, Calcium Carbonate, Monocalcium Phosphate, Magnesium Oxide, Zinc Sulfate, Manganese Sulfate, Soybean Oil, Ferrous Sulfate, Vitamin D₃ Supplement, Sulfur, Vitamin E Supplement, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydriodide, Cobalt Sulfate, Vitamin A Supplement.

FEEDING DIRECTIONS

Feed at a rate of 2.6 lbs. of feed per head per day to provide 350 mg. of Chlortetracycline per head per day. Provide fresh water and good quality hay at all times. Do not feed free choice to salt deprived or hungry cattle. Salt should be available free choice prior to using a Salt Limiting feed. Provide fresh water and good quality hay at all times.

CAUTION: Use as Directed. **Do not feed free choice to salt deprived or hungry cattle.** Keep out of reach of children. Discard empty container according to local regulations.

WARNING: Follow feeding directions. Withdraw medicated feed at least 48 hours prior to slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Do not feed to salt starved animals.

Manufactured by:

SABINE VALLEY FEEDS

Greenville, Texas 75401

Net Wt. 50 lbs. (22.7 Kg)