

SABINE VALLEY BUYRITE AP500 RANGE MEAL

#439

MEDICATED

For beef cattle on pasture.

For control of active infection of Anaplasmosis caused by *Anaplasma marginale* susceptible to Chlortetracycline. It may also be used for control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella spp.* susceptible to Chlortetracycline.

Chlortetracycline 500 grams/ton

GUARANTEED ANALYSIS

Crude Protein – (min)	14.0%
(This includes not more than 6.0% equivalent crude protein from non-protein nitrogen)	
Crude Fat – (min)	5.5%
Crude Fiber – (max).....	15.0%
Calcium – (min)	2.2%
Calcium – (max)	3.2%
Phosphorus – (min).....	0.4%
Salt – (min)	24.5%
Salt – (max)	29.5%
Potassium – (min).....	0.50%
Vitamin A – (min)	11,000 IU/LB

INGREDIENT STATEMENT

Processed Grain By-Products, Salt, Molasses, Urea, Calcium Carbonate, Magnesium Oxide, Soybean Oil, Propionic Acid, Zinc Sulfate, Manganese Sulfate, Vitamin D₃ Supplement, Copper Sulfate, Vitamin E Supplement, Potassium Sulfate, Magnesium Sulfate, Sodium Selenite, Vitamin A Supplement, Cobalt Sulfate, Ethylenediamine Dihydriodide, Amorphous Silicon Dioxide, [Citric Acid, Ethoxyquin, Butylated Hydroxytoluene, Phosphoric Acid, Butylated Hydroxyanisole as preservatives].

FEEDING DIRECTIONS

Feed to beef cattle on pasture. Feed at the rate of 1.4 pounds to provide 350 mg/head/day Chlortetracycline to control bacterial pneumonia associated with shipping fever complex caused by *Pasteurella spp.* susceptible to Chlortetracycline. In beef cattle under 700-lb. feed at the rate of 1.4 pounds/head/day to provide 350 mg/head/day Chlortetracycline to control active infection of Anaplasmosis caused by *Anaplasma marginale* susceptible to Chlortetracycline. In beef cattle over 700-lb. feed at the rate of 1 pound/head/day/500 pounds of body weight to provide 0.5 mg per pound of body weight/day Chlortetracycline to control active infection of Anaplasmosis caused by *Anaplasma marginale* susceptible to Chlortetracycline. Feed as the only source of salt and minerals. Do not feed free choice to salt deprived or hungry cattle. Salt should be available free choice prior to using a salt limiter feed. Provide fresh water and good quality hay at all times.

CAUTION: Use as Directed.

WARNING: A withdrawal period has not been established for this product in pre-ruminating calves. Do no use in calves to be processed for veal. Withdraw 48 hours prior to slaughter.

Manufactured by
SABINE VALLEY FEEDS
Greenville, Texas 75401

Net Wt. 50 lbs. (22.7 Kg)