

# SHEEP & GOAT PELLETS

400

## MEDICATED

### For Use As A Supplement For Maintenance Of Mature Breeding Animals

For the prevention of coccidiosis in young sheep caused by *Eimeria bakuensis*, *Eimeria crandallis*, *Eimeria ovinoidalis*, and *Eimeria parva*.

For the prevention of coccidiosis in young goats caused by *Eimeria christensen*, and *Eimeria ninakohlyakimovae*.

### Active Drug Ingredient

Decoquinate..... 30 gm/ton

### GUARANTEED ANALYSIS

Crude Protein – (min).....	14.0%
This feed contains not more than 2% protein equivalent from non-protein nitrogen.	
Crude Fat – (min).....	3.0%
Crude Fiber – (max).....	8.0%
Calcium – (min).....	1.8%
Calcium – (max).....	2.8%
Phosphorus – (min).....	0.8%
Salt – (min).....	0.8%
Salt – (max).....	1.8%
Copper – (min).....	7 PPM
Copper – (max).....	12 PPM
Selenium – (min).....	0.30 PPM
Vitamin A – (min).....	3,000 IU/lb.
Vitamin E – (min).....	25 IU/lb.

### INGREDIENT STATEMENT

Grain Products, Processed Grain By-Products, Plant Protein Products, Roughage Products 5%, Calcium Carbonate, Salt, Ammonium Chloride, Monocalcium Phosphate, Soybean Oil, Magnesium Oxide, Sulfur, Choline Chloride, Rice Hulls, Ascorbic Acid, Niacin Supplement, Mineral Oil, D-Calcium Pantothenate, D-Biotin, Riboflavin Supplement, Menadione Sodium Bisulfite Complex, Pyridoxine Hydrochloride, Thiamin Mononitrate, Folic Acid, Vitamin B-12 Supplement, Ferrous Sulfate, Manganese Sulfate, Zinc Sulfate, Vitamin E Supplement, Soybean Oil, Vitamin D<sub>3</sub> Supplement, Sodium Selenite, Copper Sulfate, Ethylenediamine Dihydriodide, Potassium Chloride, Vitamin A Supplement, Cobalt Sulfate, Amorphous Silicon Dioxide, [Ethoxyquin, Citric Acid, Butylated Hydroxytoluene, Phosphoric Acid, Butylated Hydroxyanisole as preservatives].

### FEEDING DIRECTIONS

**Feeding Directions:** Feed to sheep and goats as a supplement to native range. Adequate roughage should be available at all times. Good, clean water should be available at all times. Each pound of feed contains 15 mg of decoquinate. Feed 1.5 pounds of feed per 100 lbs. body to provide 22.7 mg per 100 lbs. (0.5 mg/kg) of body weight per day. Feed at least 28 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

**WARNING:** Do not feed to sheep or goats producing milk for food. Contains NPN (non-protein nitrogen), feed to ruminating animals only.

**CAUTION:** Do not use in feeds containing bentonite.

Manufactured For:  
**SABINE VALLEY FEEDS**  
3001 Bois D'Arc  
Greenville, Texas 75401

Net Wt. 50 lbs. (22.6 Kg)