

SABINE VALLEY POWER + GOAT RATION

443

MEDICATED

For Growing and/or Finishing Goats

For the prevention of coccidiosis caused by Eimeria christensenii, Eimeria crandallii and ninakohlyakimovae.

ACTIVE INGREDIENT: Monensin..... 20 grams/ton

GUARANTEED ANALYSIS

Crude Protein (Min)	16%
Includes not more than 2.5% of protein equivalent from non-protein nitrogen.	
Crude Fat (Min)	4.0%
Crude Fiber (Max)	22%
Calcium (Min)	1.0%
Calcium (Max)	2.0%
Salt (Min)	1.2%
Salt (Max)	2.2%
Phosphorus (Min)	0.45%
Copper (Min)	20 PPM
Copper (Max)	40 PPM
Selenium (Min)	0.9 PPM
Vitamin A (Min)	15,000 IU/LB
With Calf Manna, Yeast Culture and Zinpro 4-Plex	

Medicated

For Goats in confinement

INGREDIENT STATEMENT

Roughage Products (42%), Forage Products, Grain Products, Processed Grain By-Products, Plant Protein Products, Salt, Dried Whey, Calcium Carbonate, Salt, Monocalcium Phosphate, Dicalcium Phosphate, Magnesium Oxide, Zinc Oxide, Copper Sulfate, Copper Oxide, Iron Oxide, Ferrous Sulfate, Ferrous Carbonate, Manganous Oxide, Sulfur, Sodium Selenite, Vegetable Oil, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D₃ Supplement, Vitamin E Supplement, Choline Chloride, Thiamine Mononitrate, Niacin Supplement, d-Calcium Pantothenate, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B-12 Supplement, Folic Acid, Biotin, Urea, Zinc Methionine Complex, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Zinc Sulfate, Manganese Sulfate, Potassium Chloride, Cobalt Sulfate, Selenium Yeast, Dried Lactobacillus acidophilus fermentation product, Dried Enterococcus faecium fermentation product, Dextrose, Sodium Aluminosilicate, Beta Glucan (encapsulating agent), Cane Molasses, Dried Aspergillus oryzae fermentation product, Propionic Acid and/or Acetic Acid and/or Fumaric Acid and/or Formic Acid (as a preservative).

FEEDING DIRECTIONS

Feed at the rate of 2.27 lbs. per 100 lbs. of body weight to provide 22.7 mg of monensin per 100 lbs. Fresh water, good quality hay and minerals should be available at all times.

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Do not feed to lactating goats.

Manufactured for

SABINE VALLEY FEEDS

3001 Bois D' Arc, Greenville, Texas

Net Wt. 50 lbs. (22.7 Kg),